

25. J. Melillo *et al.*, in preparation.
26. GLOBALVIEW, *Cooperative Atmospheric Data Integration Project—Carbon Dioxide*. DC-ROM (National Oceanic and Atmospheric Administration/Climate Monitoring and Diagnostics Laboratory, Boulder, CO, 1998) (available via FTP at ftp.cmdl.noaa.gov, path: ccg/co2/GLOBALVIEW).
27. J. D. Mahlman, W. J. Moxim, *J. Atmos. Sci.* **35**, 1340 (1978); H. Levy, J. D. Mahlman, W. J. Moxim, *J. Geophys. Res.* **87**, 3061 (1982).
28. M. Heimann, *The Global Atmospheric Tracer Model TM2* (Max Plank Institut für Meteorologie, Hamburg, Germany, 1995).
29. M. Gloor, S.-M. Fan, S. W. Pacala, J. L. Sarmiento, M. Ramonet, *J. Geophys. Res.* **104**, 14245 (1999).
30. This research grew out of a workshop in March 1999 at Princeton University supported by the National

Oceanic and Atmospheric Administration Office of Global Programs. Additional support came from the NSF-supported National Center for Ecological Analysis and Synthesis in Santa Barbara, through its support of a working group on the carbon budget of Eurasia and North America.

9 November 2000; accepted 29 May 2001

Changes in Forest Biomass Carbon Storage in China Between 1949 and 1998

Jingyun Fang,^{1*} Anping Chen,¹ Changhui Peng,² Shuqing Zhao,¹ Longjun Ci³

The location and mechanisms responsible for the carbon sink in northern mid-latitude lands are uncertain. Here, we used an improved estimation method of forest biomass and a 50-year national forest resource inventory in China to estimate changes in the storage of living biomass between 1949 and 1998. Our results suggest that Chinese forests released about 0.68 petagram of carbon between 1949 and 1980, for an annual emission rate of 0.022 petagram of carbon. Carbon storage increased significantly after the late 1970s from 4.38 to 4.75 petagram of carbon by 1998, for a mean accumulation rate of 0.021 petagram of carbon per year, mainly due to forest expansion and regrowth. Since the mid-1970s, planted forests (afforestation and reforestation) have sequestered 0.45 petagram of carbon, and their average carbon density increased from 15.3 to 31.1 megagrams per hectare, while natural forests have lost an additional 0.14 petagram of carbon, suggesting that carbon sequestration through forest management practices addressed in the Kyoto Protocol could help offset industrial carbon dioxide emissions.

Recent studies have shown that the mid- and high-latitude forests in the Northern Hemisphere are functioning as a significant sink for C (1–4). These findings have been confirmed by several studies, mainly from North America and European countries using forest inventories (5–8). A long history of agricultural exploitation, forest management practices, and changing land use and forestry policies suggest that China, too, plays an important role in the global C cycle (9, 10). China has 133.7 million hectares of forested land (11) that range from tropical forests in the south to boreal forests in the north. Nationwide afforestation and reforestation programs have been in effect since the 1970s. To reduce the uncertainty in estimating C sinks, well-designed and statistically sound national

forest inventories over the long term, combined with direct field measurements of C stocks from local sample plots, may provide the best data sources for accurately quantifying C sinks and their dynamics at large scales.

Here, we used the National Forest Resource Inventory database for China collected from 1949 to 1998 for 5- to 10-year periods (11, 12) and a forest biomass database obtained from direct field measurements (13, 14) to estimate forest biomass C storage and its spatiotemporal distributions. The forest inventory database is based on the Forest Resource Inventory of China (FRIC), which spans seven periods: 1949, 1950–62, 1973–76, 1977–81, 1984–88, 1989–93, and 1994–98 (11, 12, 15). These inventories, excluding FRIC from 1949 which was derived from an assessment report (11, 16), were compiled from more than 250,000 plots (160,000 permanent sample plots plus 90,000 temporary sample plots) across the country. Systematic sampling with a grid of 2 km by 2 km or 4 km by 4 km and an area of 10 m by 10 m was used depending on forest region. Forest area and timber volume by age class as well as by forest type were documented at provincial levels. Unfortunately, these forest inventories do not provide detailed information about forest biomass; only the commercial portion

(such as timber volume) is available. To use timber data to estimate all forest biomass, a biomass expansion factor (BEF, defined as the ratio of all stand biomass to growing stock volume), which converts timber volume to mass and accounts for noncommercial components, such as branches, roots, and leaves, must be calculated. The forest biomass database obtained from direct field measurements (17) was used to determine BEF values for each forest type using a literature review of forest biomass studies in China (14).

Recent studies (8, 13, 18, 19) suggest that BEF is not constant, but varies with forest age, site class, stand density, and other biotic and abiotic factors that are closely associated with relative stand density, and can be expressed as a function of timber volume. Here, we used a function expressed as $BEF = a + b/x$, to obtain a variable BEF value for each forest type, where x is timber volume and a and b are constants for a forest type (13) (Table 1). Using the method published by Fang *et al.* (13), forest inventory data, and parameters listed in Table 1, forest biomass (including all living trees and shrubs) for each forest type was calculated at both provincial and national levels for all seven periods (20).

Total forest biomass C (Table 2) decreased from 5.06 Pg of C (Pg C) in 1949 to 4.38 Pg C in 1977–81, and then increased by 4.75 Pg C over the period 1980–98, mainly due to changes in land use, population growth, and economic policy changes. Since the new social system was established in 1949, rapidly increasing population and economic development have resulted in increased forest exploitation across the country (11). By 1949, Chinese forests accumulated the largest C storage (5.06 Pg C) and area-weighted mean C density (49.45 Mg ha⁻¹), due to a larger area of primary forests that have high biomass density. Since then, forest C storage has significantly decreased by 0.68 Pg C, with a mean rate of 0.022 Pg C year⁻¹ (ranging from 0.01 to 0.04 Pg C year⁻¹) from 1949 to the end of the 1970s. For this period, the policy of forest exploitation led to soil erosion, widespread desertification, loss of biodiversity, land degradation, and catastrophic flooding (21). Since the 1970s, however, the Chinese government has implemented several ecological restoration projects, including the Three-North Protective Forest Program, South China Timber Production Program, Rivers Pro-

¹Department of Urban and Environmental Science, and Key Laboratory for Earth Surface Processes of the Ministry of Education, Peking University, Beijing 100871, China. ²Ministry of Natural Resources, Ontario Forest Research Institute, 1235 Queen Street East, Sault Ste. Marie, ON, P6A 2E5, Canada, and Faculty of Forestry and The Forest Environment, Lakehead University, 955 Oliver Road, Thunder Bay, ON, P7B 5E1, Canada. ³Institute of Forest Ecology and Environment, Chinese Academy of Forestry, Beijing 100091, China.

*To whom correspondence should be addressed. E-mail: jyfang@urban.pku.edu.cn


REPORTS

tection Forest Program (22), and the Natural Forest Conservation Program (21). Consequently, forests in China have accumulated about 0.4 Pg C with an annual mean rate of 0.021 Pg C (ranging from 0.011 to 0.035 Pg C) during the past two decades (Table 2). The increasing C accumulation over this period is primarily the result of increased afforestation and reforestation (Table 2). Areas planted through afforestation and reforestation programs have expanded from 12.74×10^6 ha for the period 1970–80 to 23.11×10^6 ha in 1994–98, and forest C sequestration has increased from 0.27 to 0.72 Pg C, with a total C uptake of 0.45 Pg and a mean increased rate of 0.020 Pg C year⁻¹ (Table 2) during this time. Forest C density during this period has increased linearly from 15.32 Mg C ha⁻¹ for 1973–76 to 31.11 Mg C ha⁻¹ during the 1994–98 period, with a mean increased C uptake of 0.72 Mg ha⁻¹ year⁻¹.

To compare the contribution of planted and natural forests to C accumulation or loss, we also present total area and C content of natural forests (Table 2), which are the differences between those of all forests and planted forests since the mid-1970s. No clear changes in the area (ranging from 82.70×10^6 to 87.30×10^6 ha) and C storage (ranging from 3.9 to 4.2 Pg C) of natural forests were evident during this period, compared to planted forests. Because over this period all forests and planted forests have sequestered 0.31 Pg C (Table 2) and 0.45 Pg C (Table 2), respectively, natural forests have lost an additional 0.14 Pg C. However, natural forests have become a small C sink of 0.1 Pg during the past decade (Table 2).

There is considerable spatial (or regional) and temporal variability in forest C storage and C densities in China (Fig. 1, A and B). About 28 to 35% of forest C storage occurs in the southwestern region (including the provinces of Sichuan, Tibet, Yunnan, Guangxi, and Guizhou), followed by 24 to 31% in the northeastern region (e.g., Helongjiang, Jilin, and Liaoning provinc-

A Total forest biomass carbon storage



B Carbon density




Fig. 1. Spatiotemporal distribution of (A) total forest biomass C storage and (B) C density in China. The numbers 1 through 7 represent the periods 1949, 1950–62, 1973–76, 1977–81, 1984–88, 1989–93, and 1994–98, respectively.

Table 1. Parameters used to calculate biomass expansion factor (BEF). BEF is expressed as a function of stand timber volume (x), $BEF = a + b/x$, where a and b are constants for a forest type. Data are based on direct field measurements.

Forest type	a (Mg m ⁻³)	b (Mg)	N	R^2
<i>Abies</i> and <i>Picea</i>	0.4642	47.4990	13	0.98
<i>Betula</i>	1.0687	10.2370	9	0.70
<i>Casuarina</i>	0.7441	3.2377	10	0.95
<i>Cunninghamia lanceolata</i>	0.3999	22.5410	56	0.95
<i>Cypress</i>	0.6129	46.1451	11	0.96
Deciduous oaks	1.1453	8.5473	12	0.98
<i>Eucalyptus</i>	0.8873	4.5539	20	0.80
<i>Larix</i>	0.6096	33.8060	34	0.82
Lucidophyllous forests	1.0357	8.0591	17	0.89
Mixed conifer and deciduous forests	0.8136	18.4660	10	0.99
Mixed deciduous and <i>Sassafras</i>	0.6255	91.0013	19	0.86
Nonmerchantable woods	0.7564	8.3103	11	0.98
<i>Pinus armandii</i>	0.5856	18.7435	9	0.91
<i>P. koraiensis</i>	0.5185	18.2200	17	0.90
<i>P. massoniana</i> , <i>P. yunnanensis</i>	0.5101	1.0451	12	0.92
<i>P. sylvestris</i> var. <i>mongolica</i>	1.0945	2.0040	11	0.98
<i>P. tabulaeformis</i>	0.7554	5.0928	82	0.96
Other pines and conifer forests	0.5168	33.2378	16	0.94
<i>Populus</i>	0.4754	30.6034	10	0.87
<i>Tsuga</i> , <i>Cryptomeria</i> , <i>Keteleeria</i>	0.4158	41.3318	21	0.89
Tropical forests	0.7975	0.4204	18	0.87

Table 2. Area, total C, and C accumulation rate for all forests, and planted and natural forests for seven periods from 1949 to 1998 in China. The variables of natural forests were differences between those of all forests and planted forests.

Period	All forests				Planted forests				Natural forests		
	Total area (10 ⁶ ha)	Total C (Pg C)	C density (mg ha ⁻¹)	C accumu- lation rate (Pg C yr ⁻¹)	Total area (10 ⁶ ha)	Total C (Pg C)	C density (Mg ha ⁻¹)	C accumu- lation rate (Pg C yr ⁻¹)	Total area (10 ⁶ ha)	Total C (Pg C)	C accumu- lation rate (Pg C yr ⁻¹)
1949	102.34	5.06	49.45	–	–	–	–	–	–	–	–
1950–62	98.08	4.58	46.67	–0.040	–	–	–	–	–	–	–
1973–76	101.26	4.44	43.83	–0.010	17.39	0.27	15.32	–	83.87	4.17	–
1977–81	95.62	4.38	45.75	–0.013	12.74	0.33	25.55	0.012	82.88	4.05	–0.024
1984–88	102.19	4.45	43.53	0.011	18.74	0.52	27.48	0.027	83.45	3.93	–0.017
1989–93	108.63	4.63	42.58	0.035	21.37	0.61	28.69	0.020	87.26	4.02	0.018
1994–98	105.82	4.75	44.91	0.026	23.11	0.72	31.11	0.021	82.71	4.03	0.005

Carbon content is converted from biomass using a factor of 0.5. No information on planted forests is available before the mid-1970s. Data from Taiwan were not included in this study. Dashes indicate data that were not collected.

REPORTS

es). As expected, eastern and southeastern regions that have higher population densities and northern and northwestern regions that are arid have only a small portion of the C pool and low C density. In some regions, spatial patterns of C storage and density showed a large temporal variation. For example, forest C storage in the northwestern and northern regions has significantly increased (Fig. 1A), probably as a result of increased area afforested through the Three-North Protective Forest Program (22). Similar to C storage, higher C densities (e.g., 50 to 70 Mg ha⁻¹) were also found in the southwest and northeast of China, which are dominated by boreal forests or subalpine coniferous forests (including *Picea*, *Abies*, and *Larix*) with high biomass production. As expected, forests in the central and eastern regions, with a high proportion of plantations (31 to 40%) have lower C densities (e.g., 23 to 35 Mg ha⁻¹). This may reflect differing stand age factors (23) or perhaps greater human disturbance in these regions, which have dense population and are undergoing rapid economic development.

Although the mid- and high-latitude forests in the Northern Hemisphere greatly contribute to the terrestrial C sink, the magnitude of forest C sinks and their spatial distributions are still controversial (2, 24–26). Continental-scale comparison of C fluxes estimated using an atmospheric inverse model, terrestrial ecosystem models, and forest inventories often provide conflicting results. For example, analyses of the North American terrestrial C sink have indicated large annual variations, ranging from 0.05 to 0.17 (25) or 0.08 (27) to 1.7 Pg (2). Differences in data sources or parameters used for model simulation and estimation at different temporal and spatial scales are likely sources of uncertainty (25, 28). A narrow range of annual U.S. C sink from 0.08 to 0.35 Pg, which was estimated using inventory records (4, 6), indicates that inventory-based estimation of forest C at both regional and national scales may help to reduce the uncertainties in accurately evaluating the role of forests in regional and global C budgets. Our results, derived from 50-year monitoring at a national level, show that during the past two decades there is a mean C uptake of 0.021 Pg C year⁻¹ by Chinese forests, planted forests have accumulated 0.72 Pg total C, and their C densities have increased by about 0.72 Mg C ha⁻¹ year⁻¹ (29). The estimated C sink in China is comparable to that of North American forests, because the area and total C pool of Chinese forests are only half and one-third of those of the conterminous United States, respectively (30).

In addition, our results provide evidence

to support the proposal addressed by the Kyoto Protocol that C sequestered by afforestation or reforestation could partly offset CO₂ emissions from fossil fuel consumption, even though the increased C uptake is viewed as temporary reservoirs (31, 32). The Kyoto Protocol does not require commitments from developing countries, including China, but recent decreases in the rate of deforestation in China have already contributed to reduced CO₂ emissions (Table 2). We believe that continuing the practice of nationwide afforestation and reforestation projects could contribute significantly to global terrestrial C sinks.

References and Notes

- R. K. Dixon *et al.*, *Science* **263**, 185 (1994).
- S. Fan *et al.*, *Science* **282**, 442 (1998).
- S. C. Wofsy *et al.*, *Science* **260**, 1314 (1993).
- R. A. Houghton, J. L. Hackler, K. T. Lawrence, *Science* **285**, 574 (1999).
- E. Kauppi, K. Mielikainen, K. Kuusela, *Science* **256**, 70 (1992).
- R. A. Birdsey, L. S. Heath, in *Climate Change and the Productivity of America's Forests*, L. A. Joyce, Ed. (USDA Forest Service General Technical Report RM-228, Fort Collins, CO, 1995), pp. 1–164.
- D. P. Turner, G. J. Koerber, M. E. Harmon, J. J. Lee, *Ecol. Appl.* **5**, 421 (1995).
- S. L. Brown, P. E. Schroeder, *Ecol. Appl.* **9**, 968 (1999).
- J. Y. Fang, G. H. Liu, S. L. Xu, in *Studies on Emissions and their Mechanisms of Greenhouse Gases in China*, G. C. Wang, Y. P. Wen Eds. (China Environment Science Publishers, Beijing, 1996), pp. 81–149.
- C. H. Peng, M. J. Apps, *Tellus B* **49**, 393 (1997).
- Department of Forest Resource and Management, Chinese Ministry of Forestry, *Forest Resources of China 1949–93* (Department of Forest Resource and Management, Chinese Ministry of Forestry, Beijing, 1996).
- Chinese Ministry of Forestry, *Forest Resource Statistics of China for Periods 1950–62, 1973–76, 1977–81, 1984–88, 1989–93, and 1994–98* (Department of Forest Resource and Management, Chinese Ministry of Forestry, Beijing, 1965, 1977, 1982, 1989, 1994, and 1999).
- J. Y. Fang, G. G. Wang, G. H. Liu, S. L. Xu, *Ecol. Appl.* **8**, 1084 (1998).
- J. Y. Fang, A. P. Chen, *Forest Biomass Database of China, Technical Report* (The Center for Ecological Research & Education, Peking University, Beijing, 2000), pp. 1–50.
- In China, both the definition and timber volume measurement of forests are based on technical standards of the Forestry Ministry of China [Forestry Ministry of China, *Technical Standards for Forest Resource Measurement* (China Forestry Publishing Press, Beijing, 1983)]. According to these standards, forests are divided into natural, secondary, and planted forests, all of which have more than 30% coverage; timber volume measurements are conducted as follows: all trees with a diameter at breast height (1.3 m) (DBH) > 4 cm are measured in every sample plot, and stand timber volume is calculated by adding the volume of all individual trees within a plot.
- Forest inventory data for 1949 were not obtained from a direct survey across the country, but were derived from estimates of forest area and timber volume for major forest types for all provinces, based on forest resource statistics and forest timber records obtained from some regions, and literature reviews conducted by the Forestry Ministry of China (1996) (17).
- These field measurements provided biotic factors [stand types, species composition, stand age, stand density, tree height, DBH, basal area, stem volume, and biomass (dry mass) of different tree components], abiotic factors (e.g., climatic variables, soil type, and topography), localities (place name, latitude, longitude, and altitude), and the size of sample plots. Stem volume for individual trees and stand volume were calculated in the same way as in the forest inventory survey.
- S. Brown, A. E. Lugo, *Interciencia* **17**, 8 (1992).
- P. Schroeder, S. Brown, J. Mo, R. Birdsey, C. Cieszewski, *For. Sci.* **43**, 424 (1997).
- Based on the function $BEF = a + b/x$ in Table 1 and forest inventory data, total forest biomass (Y) of each forest type was calculated for each age class ($k = 1, 2, 3, 4, 5$), site class ($j = 1, 2, 3$), and province ($i = 1, 2, \dots, 31$) from mean stand timber volume (x_{ijk}) and its area (A_{ijk}).

$$Y = \sum_{i=1}^{31} \sum_{j=1}^3 \sum_{k=1}^5 A_{ijk} \cdot BEF \cdot x_{ijk}$$

The area-weighted mean biomass (y) of each forest type was calculated using the equation

$$y = \frac{1}{A} \sum_{i=1}^{31} \sum_{j=1}^3 \sum_{k=1}^5 A_{ijk} \cdot BEF \cdot x_{ijk}$$

where A is the total area of each forest type.

- P. Zhang *et al.*, *Science* **288**, 2135 (2000).
- The Three-North Protective Forest Program: This program started in 1978 and covers an area of 410×10^4 km², 41% of the total area of the country, across from the western part of the northeast region, the northern area of north China, and most of the northwest region (thus, Three-North) where annual precipitation is generally less than 400 mm. The Rivers Protective Forest Project: At the end of the 1980s, China has reforested along its main rivers, including the middle and upper reaches of the Yangtze River, the middle of the Yellow River, the Zuhai River valley, and the Liaohe River valley. The South China Timber Program: Mainly funded by the World Bank, China has begun to implement a large afforestation plan in South China since the mid-1980s, including the provinces of Hunan, Jiangxi, Zhejiang, Jiangsu, Fujian, Guangdong, and Hainan, to meet increasing timber demand.
- W. A. Kurz, M. J. Apps, *Ecol. Appl.* **9**, 526 (1999).
- C. B. Field, I. Y. Fung, *Science* **285**, 544 (1999).
- E. A. Holland *et al.*, *Science* **283**, 1815a (1999).
- R. A. Houghton, J. L. Hackler, *Global Ecol. Biogeogr.* **9**, 125 (2000).
- D. Schimel *et al.*, *Science* **287**, 2004 (2000).
- E.-D. Schulze *et al.*, *Science* **289**, 2058 (2000).
- Current estimates of C sinks in Chinese forests may be underestimated slightly because forest inventories are of net removals. Some removals go into long-lived products (e.g., furniture, housing, and building materials) removed from forest ecosystems. However, removals were estimated to be <2 to 3% of total biomass stocks in China [W. H. Li, F. Li, Eds., *Research of Forest Resources in China* (China Forestry Publishing House, Beijing, 1996)].
- Forest area and total forest biomass C are 200.7×10^6 ha and 12.1 Pg C for the conterminous United States (7), and 102×10^6 to 109×10^6 ha and 4.4 to 4.8 Pg C for China, respectively, during last two decades (based on the present study).
- IGBP Terrestrial Carbon Working Group, *Science* **280**, 1393 (1998).
- B. Schlamadinger, T. Karjalainen, in *A Special Report of the IPCC: Land Use, Land-Use Change and Forestry*, R. Watson *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 2000), pp. 127–180.
- We thank S. Brown, L. Buse, and R. Houghton for their valuable comments on an earlier draft, and J. J. Chen for assistance in inventory data collection. This research was supported by the State Key Basic Research and Development Plan (G2000046801) and National Natural Science Foundation of China (39425003 and 40024101) to J.Y.F. C.H.P. acknowledges the support of the Ontario Climate Change Program of Ministry of Natural Resources.

26 December 2000; accepted 13 March 2001

Changes in Forest Biomass Carbon Storage in China Between 1949 and 1998

Jingyun Fang, Anping Chen, Changhui Peng, Shuqing Zhao and Longjun Ci

Science **292** (5525), 2320-2322.
DOI: 10.1126/science.1058629

ARTICLE TOOLS

<http://science.sciencemag.org/content/292/5525/2320>

RELATED CONTENT

<http://science.sciencemag.org/content/sci/292/5525/2261.full>
<http://science.sciencemag.org/content/sci/292/5525/2316.full>
<http://science.sciencemag.org/content/sci/296/5572/1359.full>

REFERENCES

This article cites 19 articles, 11 of which you can access for free
<http://science.sciencemag.org/content/292/5525/2320#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)